

Supersymmetric Electroweak Corrections to Top Quark Production at LHC

Hong-Yi Zhou

*CCAST (World Laboratory), P. O. Box 8730, Beijing 100080, P.R. China,
Institute of Modern Physics and Department of Physics,
Tsinghua University, Beijing 100084, P.R. China**

Chong-Sheng Li

*CCAST (World Laboratory), P. O. Box 8730, Beijing 100080, P.R. China,
Physics Department, Peking University, Beijing 100871, P.R. China**
(February 1, 2008)

Abstract

The $\alpha m_t^2/m_W^2$ order supersymmetric electroweak corrections arising from loops of chargino, neutralino, and squark to top quark pair production by gg fusion at LHC are calculated in the minimal supersymmetric model. We found that the corrections amount about a few percent.

I. Introduction

The top quark has been found experimentally by the CDF and D0 Collaborations with the mass and production cross section $m_t = 176 \pm 8(stat) \pm 10(syst)$ GeV $\sigma = 6.8^{+3.6}_{-2.4} pb$, and $m_t = 199^{+19}_{-20}(stat) \pm 22(syst)$ GeV $\sigma = 6.4 \pm 2.2 pb$, respectively [1]. Although this measured mass is close to the central value predicted by the best fit of the Standard Model (SM) to the latest LEP data, the central value of the cross section is somewhat larger than the Standard Model prediction $\sigma_{t\bar{t}} = 5.52^{+0.07}_{-0.45}$ pb for $m_t = 175$ GeV at $\sqrt{s} = 1.8$ TeV $p\bar{p}$ collider in which the effects of multiple soft-gluon emissions have been properly resummed [2]. In addition, there are still a number of unsolved theoretical problems in the SM. New physics beyond the SM are still possible. Among various models of new physics so far considered, supersymmetry (SUSY) is a promising one at present. The simplest and interesting SUSY model is the minimal supersymmetric extension of the standard model (MSSM) [3]. At the future multi-TeV proton colliders such as the CERN Large Hadron Collider (LHC), $t\bar{t}$ production will be enormously larger than the Tevatron rates and the accuracy with which the top quark production cross section can be measured will be much better (the uncertainty is about 5% at LHC [4]). Thus theoretical calculations of the radiative corrections to the production of the top quark at those colliders are of importance.

QCD corrections to $O(\alpha_s^3)$ and electroweak one-loop corrections in the SM to $t\bar{t}$ production in hadron colliders are carried out in Ref. [2] [5] and Ref. [6]. Yukawa corrections to $t\bar{t}$ production at the Fermilab Tevatron and LHC in two-Higgs-doublet models are calculated in [7] [8]. In the MSSM, electroweak corrections from chargino, neutralino and squark to the top pair production via $q\bar{q}$ annihilation at the Fermilab Tevatron are calculated in Ref. [9]. Recent calculation of the supersymmetric QCD corrections to the top quark pair production at the Tevatron shows that they increase the cross section by about 20% [10] [11]. But this is still within the experimental uncertainty 30%. At LHC the main production mechanism of top quark pair is the gluon-gluon fusion process $gg \rightarrow t\bar{t}$. In this paper we investigate the electroweak corrections of order $\alpha m_t^2/m_W^2$ arising from chargino, neutralino and squark to the top quark production by the process $gg \rightarrow t\bar{t}$ at LHC. The formalism of the calculation of the corrections to the matrix elements will be given in Sec.II. In Sec III, we present our numerical examples and discussions of the corrections to the cross sections in the MSSM.

II. Formalism

The tree-level Feynman diagrams and the relevant supersymmetric electroweak corrections to $gg \rightarrow t\bar{t}$ are shown in Fig.1 (u-channel diagrams of (b) and (e)–(h) are not explicitly shown) in which the dashed lines in the loop represent the squark $\tilde{t}_i$ or $\tilde{b}_i$ ($i = 1, 2$) and the solid lines represent neutralinos or charginos, respectively.

The supersymmetric partner of left- and right-handed massive quarks mix [12]. The mass eigenstates $\tilde{q}_1$ and $\tilde{q}_2$ are related to the current eigenstates $\tilde{q}_L$ and $\tilde{q}_R$

$$\tilde{q}_1 = \tilde{q}_L \cos \theta_q + \tilde{q}_R \sin \theta_q, \quad \tilde{q}_2 = -\tilde{q}_L \sin \theta_q + \tilde{q}_R \cos \theta_q \quad (1)$$

The mixing angle θ_t and the masses $m_{\tilde{t}_1}$, $m_{\tilde{t}_2}$ can be calculated by diagonalizing the following mass matrix

$$\begin{aligned}
M_t^2 &= \begin{pmatrix} M_{\tilde{t}_L}^2 & m_t m_{LR} \\ m_t m_{LR} & M_{\tilde{t}_R}^2 \end{pmatrix} \\
M_{\tilde{t}_L}^2 &= m_{\tilde{t}_L}^2 + m_t^2 + \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W\right) \cos(2\beta) m_Z^2 \\
M_{\tilde{t}_R}^2 &= m_{\tilde{t}_R}^2 + m_t^2 + \frac{2}{3} \sin^2 \theta_W \cos(2\beta) m_Z^2 \\
m_{LR} &= \mu \cot \beta + A_t
\end{aligned} \tag{2}$$

where $m_{\tilde{t}_L}^2$, $m_{\tilde{t}_R}^2$ are the soft SUSY-breaking mass terms of left- and right-handed stops, μ is the coefficient of the $H_1 H_2$ mixing term in the superpotential, A_t is the parameter describing the strength of soft SUSY-breaking trilinear scalar interaction $\tilde{t}_L \tilde{t}_R H_2$, $\tan \beta = v_2/v_1$ is the ratio of the vacuum expectation values of the two Higgs doublets.

From Eq.(2), we can get the expressions for $m_{\tilde{t}_{1,2}}^2$ and θ_t :

$$m_{\tilde{t}_{1,2}}^2 = \frac{1}{2} \left[M_{\tilde{t}_L}^2 + M_{\tilde{t}_R}^2 \mp \sqrt{(M_{\tilde{t}_L}^2 - M_{\tilde{t}_R}^2)^2 + 4m_t^2 m_{LR}^2} \right] \tag{3}$$

$$\tan \theta_t = \frac{m_{\tilde{t}_1}^2 - M_{\tilde{t}_L}^2}{m_t m_{LR}} \tag{4}$$

For the sbottoms, we neglect the mixing between the left- and right- handed sbottoms($\theta_b = 0$) and have

$$m_{\tilde{b}_{1,2}}^2 = m_{\tilde{t}_L, \tilde{b}_R}^2 + m_b^2 \pm (T_{L,R}^3 - Q_b \sin \theta_W) \cos(2\beta) m_Z^2, \tag{5}$$

where $T_{L,R}^3 = -\frac{1}{2}$, 0 , $Q_b = -\frac{1}{3}$ and $m_{\tilde{t}_L, \tilde{b}_R}^2$ are the soft SUSY-breaking mass terms for left- and right-handed sbottoms.

In the presence of squark mixing, the squark-quark-neutralino and squark-quark-chargino interaction Lagrangian of order gm_t/m_W is given by

$$\begin{aligned}
L_{\tilde{\chi} \tilde{q} \tilde{q}} &= -\frac{gm_t}{\sqrt{2}m_W \sin \beta} \sum_{j=1}^4 \tilde{t} [(a_{\tilde{t}_{1j}} - b_{\tilde{t}_{1j}} \gamma_5) \tilde{t}_1 + (a_{\tilde{t}_{2j}} - b_{\tilde{t}_{2j}} \gamma_5) \tilde{t}_2] \tilde{\chi}_j^0 \\
&+ \frac{gm_t}{\sqrt{2}m_W \sin \beta} \sum_{j=1}^2 \tilde{t} [(a_{\tilde{b}_{1j}} - b_{\tilde{b}_{1j}} \gamma_5) \tilde{b}_1 + (a_{\tilde{b}_{2j}} - b_{\tilde{b}_{2j}} \gamma_5) \tilde{b}_2] \tilde{\chi}_j^+ + H.C. ,
\end{aligned} \tag{6}$$

where g is the SU(2) coupling constant, and $a_{\tilde{t}_{1j}}$, $b_{\tilde{t}_{1j}}$, $a_{\tilde{t}_{2j}}$, $b_{\tilde{t}_{2j}}$, $a_{\tilde{b}_{1j}}$, $b_{\tilde{b}_{1j}}$, $a_{\tilde{b}_{2j}}$, $b_{\tilde{b}_{2j}}$ are given by

$$\begin{aligned}
a_{\tilde{t}_{1j}} &= \frac{1}{2} (N_{j4}^* \cos \theta_t + N_{j4} \sin \theta_t), \quad b_{\tilde{t}_{1j}} = \frac{1}{2} (N_{j4}^* \cos \theta_t - N_{j4} \sin \theta_t), \\
a_{\tilde{t}_{2j}} &= \frac{1}{2} (-N_{j4}^* \sin \theta_t + N_{j4} \cos \theta_t), \quad b_{\tilde{t}_{2j}} = \frac{1}{2} (-N_{j4}^* \sin \theta_t - N_{j4} \cos \theta_t), \\
a_{\tilde{b}_{1j}} &= b_{\tilde{b}_{1j}} = \frac{1}{2} V_{j2}^* \cos \theta_b, \quad a_{\tilde{b}_{2j}} = b_{\tilde{b}_{2j}} = -\frac{1}{2} V_{j2}^* \sin \theta_b
\end{aligned} \tag{7}$$

V_{j2} are the elements of 2×2 matrix V and N_{j4} are the elements of 4×4 matrix N (see the Appendix).

At the tree level, the S-matrix element is composed of three different production channels(s-,t-,u-channel) as follows:

$$\begin{aligned} M_0^s &= -ig_s^2(if_{abc}T^c)_{ji}\bar{u}(p_2)\not{p}_1 v(p_1)/\hat{s} \\ &= -i(T^a T^b - T^b T^a)_{ji} M_0^{s'}, \end{aligned} \quad (8)$$

$$\begin{aligned} M_0^t &= -ig_s^2(T^b T^a)_{ji}\bar{u}(p_2)\not{p}_4(\not{q} + m_t)\not{p}_3 v(p_1)/(\hat{t} - m_t^2) \\ &= -i(T^b T^a)_{ji} M_0^{t'}, \end{aligned} \quad (9)$$

$$\begin{aligned} M_0^u &= M_0^t(p_3 \leftrightarrow p_4, \quad T^a \leftrightarrow T^b, \quad \hat{t} \rightarrow \hat{u}) \\ &= -i(T^a T^b)_{ji} M_0^{u'}, \end{aligned} \quad (10)$$

where $q = p_2 - p_4$, $\epsilon_4^\mu = \epsilon^\mu(p_4)$, $\epsilon_3^\mu = \epsilon^\mu(p_3)$, Γ^μ is given in the Appendix. Instead of calculating the square of the amplitudes explicitly, we calculate the helicity amplitudes numerically by using the method of Ref. [13]. This method greatly simplifies our calculations.

The $O(\alpha m_t^2/m_W^2)$ SUSY electroweak corrections to $gg \rightarrow t\bar{t}$ are shown in Fig.1 (c)–Fig.1 (h). The sum of them is QCD gauge invariant without the strong coupling constant renormalization. In our calculation, we use dimensional regularization to regulate the ultraviolet divergences and adopt the on-mass-shell renormalization scheme. We also discard the terms proportional to γ_5 .

We only give the explicitly results of the s- and t-channel contributions to the SUSY electroweak corrections. The u-channel results can be obtained by the following substitutions:

$$p_3 \leftrightarrow p_4, \quad T^a \leftrightarrow T^b, \quad \hat{t} \leftrightarrow \hat{u}. \quad (11)$$

Fig.1(c) lead to the s-channel vertex correction δM^{s1} :

$$\begin{aligned} \delta M^{s1} &= -ig_s^2(if_{abc}T^c)_{ji}\bar{u}(p_2)(F_0^{s1} + F_1^{s1}\not{p}_1 + F_6^{s1})v(p_1)/\hat{s} \\ &= -i(T^a T^b - T^b T^a)_{ji} \delta M^{s1'} \end{aligned} \quad (12)$$

Fig.1(d) gives δM^{s2} :

$$\begin{aligned} \delta M^{s2} &= -ig_s^2(T^a T^b + T^b T^a)_{ji}\bar{u}(p_2)F_0^{s2}v(p_1) \\ &= -i(T^b T^a + T^a T^b)_{ji} \delta M^{s2'} \end{aligned} \quad (13)$$

The top quark self-energy $\delta M^{self,t}$ of Fig.1(e) is:

$$\begin{aligned} \delta M^{self,t} &= -ig_s^2(T^b T^a)_{ji}\bar{u}(p_2)\not{p}_4(\not{q} + m_t) \\ &\quad [F_1^{self,t} + F_2^{self,t}\not{q}](\not{q} + m_t)\not{p}_3 v(p_1)/(\hat{t} - m_t^2) \\ &= -i(T^b T^a)_{ji} \delta M^{self,t'} \end{aligned} \quad (14)$$

Vertex correction $\delta M^{v1,t}$ of Fig.1(f) is:

$$\begin{aligned} \delta M^{v1,t} &= -ig_s^2(T^b T^a)_{ji}\bar{u}(p_2)(F_0^{v1,t} + F_1^{v1,t}\not{p}_4 + F_6^{v1,t})(\not{q} + m_t)\not{p}_3 v(p_1)/(\hat{t} - m_t^2) \\ &= -i(T^b T^a)_{ji} \delta M^{v1,t'} \end{aligned} \quad (15)$$

Vertex correction $\delta M^{v2,t}$ of Fig.1(g) is:

$$\begin{aligned}\delta M^{v2,t} &= -ig_s^2(T^b T^a)_{ji}\bar{u}(p_2)\not{\epsilon}_4(\not{q} + m_t)(F_0^{v2,t} + F_1^{v2,t}\not{\epsilon}_3 + F_6^{v2,t})v(p_1)/(\hat{t} - m_t^2) \\ &= -i(T^b T^a)_{ji}\delta M^{v2,t'}\end{aligned}\quad (16)$$

$\delta M^{box,t}$ of the box diagram Fig.1(h) is:

$$\begin{aligned}\delta M^{box,t} &= -ig_s^2(T^b T^a)_{ji}\bar{u}(p_2)[F_0^{b,t} + F_3^{b,t}]v(p_1) \\ &= -i(T^b T^a)_{ji}\delta M^{box,t'}\end{aligned}\quad (17)$$

The total amplitude can be written as:

$$M_{ji} = -i[(M_0^+ + \delta M^+)O_{ji}^{(+)} + (M_0^- + \delta M^-)O_{ji}^{(-)}], \quad (18)$$

where

$$\begin{aligned}O^{(+)} &= \frac{T^b T^a + T^a T^b}{2}, \quad O^{(-)} = \frac{T^b T^a - T^a T^b}{2} \\ M_0^+ &= M_0^{t'} + M_0^{u'}, \quad M_0^- = M_0^{t'} - M_0^{u'} - 2M_0^{s'}, \\ \delta M^+ &= 2\delta M^{s2'} + \delta M^{self,t'} + \delta M^{self,u'} + \delta M^{v1,t'} + \delta M^{v1,u'} \\ &\quad + \delta M^{v2,t'} + \delta M^{v2,u'} + \delta M^{box,t'} + \delta M^{box,u'} \\ \delta M^- &= -2\delta M^{s1'} + \delta M^{self,t'} - \delta M^{self,u'} + \delta M^{v1,t'} - \delta M^{v1,u'} \\ &\quad + \delta M^{v2,t'} - \delta M^{v2,u'} + \delta M^{box,t'} - \delta M^{box,u'}\end{aligned}\quad (19)$$

The color sum of the corrected amplitude square is:

$$\begin{aligned}\sum_{color} |M|^2 &= \frac{7}{3}|M_0^+|^2 + 3|M_0^-|^2 \\ &\quad + \frac{14}{3}Re(M_0^+ \delta M^{+\dagger}) + 6Re(M_0^- \delta M^{-\dagger})\end{aligned}\quad (20)$$

The spin sum as well as phase-space integration and parton distribution convolution are done by the VEGAS program. The correction cross section $\Delta\sigma$ is defined as

$$\Delta\sigma = \sigma - \sigma_0, \quad (21)$$

where σ is the cross section given by $|M|^2$ and σ_0 is the tree level QCD cross section given by $|M_0^+|^2$ and $|M_0^-|^2$.

III. Numerical Examples and Discussions

The production cross section is obtained by convoluting the partonic cross section with certain parton distribution. The relative correction is not sensitive to parton distribution. In this paper, we take the Martin-Roberts-Stirling (MRS) parton distribution set A' [14] with $Q^2 = \hat{s}$. The following parameters are used in our calculation:

$$\begin{aligned}\sqrt{s} &= 14 \text{ TeV}, \quad m_t = 176 \text{ GeV}, \quad m_b = 4.9 \text{ GeV}, \\ m_W &= 80.22 \text{ GeV}, \quad m_Z = 91.175 \text{ GeV}, \quad \alpha = \frac{1}{128.8}.\end{aligned}\quad (22)$$

Care must be taken in the calculation of the form factors expressed in terms of the standard loop integrals defined in Ref. [15]. As has been discussed in Ref. [16], the formulae for the form factors given in terms of the tensor loop integrals will be ill-defined when the scattering is forwards or backwards wherein the Gram determinants of some matrices vanish and thus their inverses do not exist. This problem can be avoided by taking the kinematic cuts on the rapidity y and the transverse momentum p_T . In this paper, we take

$$|y| < 2.5, \quad p_T > 20 \text{ GeV}. \quad (23)$$

The cuts will also increase the relative corrections [6].

We first checked the QCD gauge invariance by the substitution $p_4 \rightarrow \epsilon_4$ and $p_3 \rightarrow \epsilon_3$ and find that δM^+ , δM^- are a few order of magnitudes smaller.

In the calculation of the chargino and neutralino masses, we fix $M = 200 \text{ GeV}$, $\mu = -100 \text{ GeV}$ and use the relation $M' = \frac{5g'^2}{3g^2}M$ (see the Appendix). We also assume $m_{\tilde{t}_L} = m_{\tilde{t}_R} = m_{\tilde{b}_R} = m_{\tilde{q}}$.

The relative correction to the hadronic cross section as a function of the squark mass parameter $m_{\tilde{q}}$ with $\tan \beta = 1$ and $m_{LR} = 0$ (corresponding to non-mixing case) is presented in Fig.2. For $\tan \beta = 1$, the chargino masses $m_{\tilde{\chi}_j^\pm} = (220, 120) \text{ GeV}$ and the neutralino masses $m_{\tilde{\chi}_j^0} = (105, 221, 128, 100) \text{ GeV}$. The correction is always negative. For $m_{\tilde{q}} < 150 \text{ GeV}$, the correction is very sensitive to $m_{\tilde{q}}$. A sharp dip at about $m_{\tilde{q}} = 56 \text{ GeV}$ is due to the singularity of the top quark wave function renormalization constant at the threshold point $m_t = m_{\tilde{b}_1} + m_{\tilde{\chi}_2^+}$ (note that for $\tan \beta = 1$, $m_{\tilde{b}_1} = m_{\tilde{q}}$). This singularity will disappear if the finite widths of the top quark and the charginos are taken into account. The correction exceeds -5% only in a small region near the dip. The correction approaches to zero at large $m_{\tilde{q}}$ which shows the decoupling behaviour.

Fig.3 shows the dependence of the relative correction to the hadronic cross section on the stop mixing parameter m_{LR} . We set $m_{\tilde{q}} = 100 \text{ GeV}$ and $\tan \beta = 1$. m_{LR} affects the mass splitting and mixing angle of $\tilde{t}_1$ and $\tilde{t}_2$. The mass splitting increases as m_{LR} increases. We fix $\tilde{t}_1$ to be the light one (cf. Eq.(3)). Therefore, $\theta_t = \frac{\pi}{4}$ for $m_{LR} < 0$, $\theta_t = -\frac{\pi}{4}$ for $m_{LR} > 0$. The mixing angle causes the asymmetry of the relative correction between $m_{LR} > 0$ and $m_{LR} < 0$ although the mass splitting of $\tilde{t}_1$ and $\tilde{t}_2$ is symmetry between $m_{LR} > 0$ and $m_{LR} < 0$. The dip at about $m_{LR} = -200 \text{ GeV}$ is due to the threshold effect at $m_t = m_{\tilde{t}_1} + m_{\tilde{\chi}_j^0}$. No dip is found at $m_{LR} = 200 \text{ GeV}$ because when $m_{LR} > 0$, $\theta_t = -\pi/4$, the $t\tilde{t}_1\tilde{\chi}_j^0$ coupling is proportional to γ_5 ($a_{ij} = 0$, $b_{ij} \neq 0$). From the expression of the top quark renormalization constant (see the Appendix), one can see that the singularities of G_0 and G_1 cancel with each other when $a_{ij} = 0$, $b_{ij} \neq 0$.

In Fig.4, we present the $\tan \beta$ dependence of the relative correction to the hadronic cross section at given $m_{\tilde{q}} = 100 \text{ GeV}$ and $m_{LR} = 100 \text{ GeV}$. $\tan \beta$ slightly affects the stop mass splitting and $m_{\tilde{b}_1}$. The factor $1/\sin^2 \beta$ in the coupling constant leads to the rapid increase of the correction in the range $\tan \beta < 1$. But the increase is somewhat more quickly than $1/\sin^2 \beta$ because $m_{\tilde{b}_1}$ decreases as β decreases.

From Fig.2-4, we see that only for $\tan \beta < 1$ and a small region near the threshold $t \rightarrow \tilde{b}_1\tilde{\chi}_j^+$ and $t \rightarrow \tilde{t}_1\tilde{\chi}_j^0$ the correction may exceed -5%. Otherwise, the correction amounts only a few percent smaller than -5%. Therefore, we conclude that the supersymmetric electroweak corrections of order $\alpha m_t^2/m_W^2$ to top quark pair production at LHC are potentially observable for $\tan \beta < 1$ and small parameter region near the threshold $t \rightarrow \tilde{b}_1\tilde{\chi}_j^+$, $t \rightarrow \tilde{t}_1\tilde{\chi}_j^0$.

ACKNOWLEDGMENTS

This work is supported in part by the National Natural Science Foundation of China, the Fundamental Research Foundation of Tsinghua University and a grant from the state commission of Science and Technology of China.

Appendix

We give here the form factors for the matrix element appeared in the text. They are written in terms of the conventional one-, two-, three- and four-point scalar loop integrals defined in Ref. [15].

$$\begin{aligned}
F_0^{s1} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[m_j(a_{ij}^2 - b_{ij}^2)((p_2 - p_1) \cdot \Gamma C_0 \\
&\quad - 2C^{10} \cdot \Gamma)](-p_2, k, m_j, m_i, m_i) \\
F_1^{s1} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[\delta Z_{ij}^v] \\
F_6^{s1\mu} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[(a_{ij}^2 + b_{ij}^2)((p_2 - p_1) \cdot \Gamma C^{10\mu} \\
&\quad - 2C^{21\mu}(\Gamma))](-p_2, k, m_j, m_i, m_i) \\
F_0^{s2} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[m_t(a_{ij}^2 + b_{ij}^2)C_{11} - m_j(a_{ij}^2 - b_{ij}^2)C_0](-p_2, k, m_j, m_i, m_i) \\
F_1^{self,t} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[-m_j(a_{ij}^2 - b_{ij}^2)B_0 + \delta m_{ij} + m_t \delta Z_{ij}^v](\hat{t}, m_j^2, m_i^2)/(\hat{t} - m_t^2) \\
F_2^{self,t} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[(a_{ij}^2 + b_{ij}^2)B_1 - \delta Z_{ij}^v](\hat{t}, m_j^2, m_i^2)/(\hat{t} - m_t^2) \\
F_0^{v1,t} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[2m_j(a_{ij}^2 - b_{ij}^2)(p_2 \cdot \epsilon_4 C_0 - C^{10} \cdot \epsilon_4)](-p_2, p_4, m_j, m_i, m_i) \\
F_1^{v1,t} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[\delta Z_{ij}^v] \\
F_6^{v1,t\mu} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[2(a_{ij}^2 + b_{ij}^2)(p_2 \cdot \epsilon_4 C^{10\mu} \\
&\quad - C^{21\mu}(\epsilon_4))](-p_2, p_4, m_j, m_i, m_i) \\
F_0^{v2,t} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[2m_j(a_{ij}^2 - b_{ij}^2)(-p_1 \cdot \epsilon_3 C_0 \\
&\quad - C^{10} \cdot \epsilon_3)](p_1, -p_3, m_j, m_i, m_i) \\
F_1^{v2,t} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[\delta Z_{ij}^v] \\
F_6^{v2,t\mu} &= - \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[2(a_{ij}^2 + b_{ij}^2)(-p_1 \cdot \epsilon_3 C^{10\mu} \\
&\quad - C^{21\mu}(\epsilon_3))](p_1, -p_3, m_j, m_i, m_i) \\
F_0^{b,t} &= \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[4m_j(a_{ij}^2 - b_{ij}^2)(p_2 \cdot \epsilon_4 p_1 \cdot \epsilon_3 D_0 - p_1 \cdot \epsilon_3 D^{10} \cdot \epsilon_4 \\
&\quad + p_2 \cdot \epsilon_4 D^{10} \cdot \epsilon_3 - \epsilon_3 \cdot D^{21}(\epsilon_4))](-p_2, p_4, p_3, m_j, m_i, m_i, m_i)
\end{aligned}$$

$$F_3^{b,t\mu} = \sum_j \sum_{i=\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2} CPL[4(a_{ij}^2 + b_{ij}^2)(p_2 \cdot \epsilon_4 p_1 \cdot \epsilon_3 D^{10\mu} - p_1 \cdot \epsilon_3 D^{21\mu}(\epsilon_4) \\ + p_2 \cdot \epsilon_4 D^{21\mu}(\epsilon_3) - D^{32\mu}(\epsilon_3, \epsilon_4))(-p_2, p_4, p_3, m_j, m_i, m_i, m_i)]$$

In the above i, j summation, $j = 1, 2, 3, 4$, $m_j = m_{\tilde{\chi}_j^0}$ for $i = \tilde{t}_1, \tilde{t}_2$, $j = 1, 2$, $m_j = m_{\tilde{\chi}_j^+}$ for $i = \tilde{b}_1, \tilde{b}_2$. As $\theta_b = 0$, $\tilde{b}_2$ actually does not contribute to the sum(cf. Eq.(7)).

$$k = p_1 + p_2 = p_3 + p_4, \quad \hat{s} = k^2, \quad \hat{t} = q^2 = (p_2 - p_4)^2, \quad \hat{u} = (p_2 - p_3)^2, \\ \Gamma^\mu = (-p_4 + p_3)^\mu \epsilon_3 \cdot \epsilon_4 + (2p_4 + p_3) \cdot \epsilon_3 \epsilon_4^\mu - (2p_3 + p_4) \cdot \epsilon_4 \epsilon_3^\mu, \\ CPL = \frac{\alpha m_t^2}{8\pi m_W^2 \sin^2 \theta_W \sin^2 \beta}$$

$$C^{10\mu} = C^\mu, \quad C^{21\mu}(p) = p_\nu C^{\mu\nu}, \quad C^{20} = g_{\mu\nu} C^{\mu\nu} - \frac{1}{2}, \\ D^{10\mu} = D^\mu, \quad D^{21\mu}(p) = p_\nu D^{\mu\nu}, \quad D^{20} = g_{\mu\nu} D^{\mu\nu}, \\ D^{32\mu}(p, l) = p_\nu l_\alpha D^{\mu\nu\alpha}, \quad D^{30\mu} = g_{\nu\alpha} D^{\mu\nu\alpha}.$$

In our calculation, we calculate the tensor loop integrals C^μ , $C^{\mu\nu}$, D^μ , $D^{\mu\nu}$ and $D^{\mu\nu\alpha}$ numerically instead of expanding them explicitly.

The renormalization constants are:

$$Z_{ij}^v = [(a_{ij}^2 + b_{ij}^2)(B_1 + 2m_t^2 G_1) + 2(a_{ij}^2 - b_{ij}^2)m_t m_j G_0](p^2, m_j^2, m_i^2)|_{p^2=m_t^2} \\ \delta m_{ij} = [-(a_{ij}^2 + b_{ij}^2)m_t B_1 + (a_{ij}^2 - b_{ij}^2)m_j B_0](p^2, m_j^2, m_i^2)|_{p^2=m_t^2}$$

$$\text{where } G_0 = -\frac{\partial B_0(p^2, m_i', m_i)}{\partial p^2}, \quad G_1 = \frac{\partial B_1(p^2, m_i', m_i)}{\partial p^2}.$$

The neutralino and chargino masses, the 2×2 matrix V and the 4×4 matrix N are given by the following relations [3]:

$$X = \begin{pmatrix} M & m_W \sqrt{2} \sin \beta \\ m_W \sqrt{2} \cos \beta & \mu \end{pmatrix},$$

$$V X^2 V^{-1} = M_D^2$$

$$Y = \begin{pmatrix} M' & 0 & -m_Z s_W \cos \beta & m_Z s_W \sin \beta \\ 0 & M & m_Z c_W \cos \beta & -m_Z c_W \sin \beta \\ -m_Z s_W \cos \beta & m_Z c_W \cos \beta & 0 & -\mu \\ m_Z s_W \sin \beta & -m_Z c_W \sin \beta & -\mu & 0 \end{pmatrix},$$

$$N Y^2 N^{-1} = N_D^2, \quad N^* Y N^{-1} = N_D,$$

where M, M' are the masses of gauginos corresponding to $SU(2)$ and $U(1)$, respectively. With the grand unification assumption, we have the relation $M' = \frac{5}{3}(g'^2/g^2)M$. M_D^2 is a diagonal matrix and the chargino mass squares correspond to its diagonal elements. N_D is a diagonal matrix with non-negative entries which give the masses of neutralinos. The chargino and neutralino masses depend on the parameters M, μ and $\tan \beta$.

REFERENCES

- [1] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74** (1995) 2626; D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **74** (1995) 2632.
- [2] E.L. Berger and H. Contopanagos, ANL-HEP-PR-95-31, May 1995; see also: E.Laenen, J.Smith, W.L. van Neeven, Nucl. Phys. **B369**(1992) 543; Phys. Lett. **B321**(1994) 254.
- [3] H.E.Haber and G.L.Kane, Phys. Rep. **117**(1985) 75; J.F.Gunion and H.E.Haber, Nucl.Phys. **B272** (1986) 1; Erratum, Nucl. Phys. **B402** (1993) 567.
- [4] T.M.Liss, hep-ph/9501274.
- [5] P. Nason, S.Dawson and R.K.Eliss, Nucl. Phys.**B303** (1988) 607; G.Altarelli, M. Diemoz, G.Martinelli and P. Nason, Nucl. Phys.**B308** (1988) 724; W.Beenakker, H.Kuijff, W.L.van Neerven and J.Smith, Phys. Rev. **D40** (1989) 54.
- [6] W. Beenakker, A.Denner, W.Hollik, R.Mertig, T.Sack and D. Wackeroth, Nucl. Phys. **B411** (1994) 343.
- [7] A. Stange and S. Willenbrock, Phys. Rev. **D48** (1993) 2054.
- [8] H.Y. Zhou, C.S. Li and Y.P.Kuang, hep-ph/9603435.
- [9] J.M.Yang and C.S. Li, Phys. Rev. **D52** (1995) 1541; J.M.Yang and C.S. Li, hep-ph/9603442.
- [10] C.S. Li,B.Q. Hu, J.M. Yang and C.G. Hu, Phys. Rev. **D52** (1995) 5014; C.S. Li, H.Y. Zhou, Y.L. Zhu and J.M. Yang, Phys. Lett. **B379** (1996) 135.
- [11] J.Kim, J.L. Lopez, D.V. Nanopoulos and R. Rangarajan, Phys. Rev. **D 54**,4364 (1996).
- [12] J.Ellis and S.Rudaz, Phys. Lett. **B128** (1983) 248; A.Bouquet, J.Kaplan and C.Savoy, Nucl. Phys. **B262** (1985) 299.
- [13] K.Hagiwara and D.Zeppenfeld, Nucl. Phys. **B313** (1989) 560; V.Barger, T.Han, and D.Zeppenfeld, Phys.Rev. **D41** (1990) 2782.
- [14] A.D. Martin, W.J. Stirling and R.G. Roberts, Phys. Lett. **B354** (1995) 155.
- [15] G. Passarino and M.Veltman, Nucl. Phys. **B160** (1979) 151; G.'t Hooft and M. Veltman, Nucl. Phys. **B153** (1979) 365.
- [16] A. Denner, Fortschr. Phys. **41**,No.4 (1993) 307.

Figure Captions

Fig.1 Feynman diagrams of tree level and $O(\alpha m_t^2/m_W^2)$ SUSY electroweak correction.

Fig.2 Relative correction to hadronic cross section versus $m_{\tilde{q}}$ with $\tan\beta = 1$ and $m_{LR} = 0$.

Fig.3 Relative correction to hadronic cross section versus m_{LR} with $\tan\beta = 1$ and $m_{\tilde{q}} = 100$ GeV.

Fig.4 Relative correction to hadronic cross section versus $\tan\beta$ with $m_{\tilde{q}} = 100$ GeV and $m_{LR} = 100$ GeV.




Fig. 1




Fig. 2